

## ***Allium oleraceum* L.**

## **Field Garlic**

Native

GB: Vulnerable

England: Least Concern

VC5 Rare; VC6 Not Scarce


A bulbous perennial of dry field edges, hedgerows, lane sides and other grassy places. In VC5 this species was first recorded in 1911, when one plant was seen on the bank of a lane south of Staple Fitzpaine (Marshall, 1914). In 1924 it was found on Wembdon Hill by E.J. Hamlin; there have been no further records for these two sites. In 1975, C.A. Howe found this species at Cleeve Hill, which was the only VC5 site given by Green *et al.* (1997). Since then, it has been found at Ford Farm on Exmoor, Old Cleeve, Huish and Currymead Lane, south of Curry Rivel. In VC6 *Allium oleraceum* is not scarce but has a distribution largely restricted to two areas. It is particularly found around the Poldens, southeast to Lytes Cary, and also south of the River Avon near Bristol, from Pill to Leighwoods. It was discovered at Tyntesfield (NT estate) in 2004 by Pam Millman. In 2011, Rob Randall found a few plants near Southstoke, where it was collected in 1851 by L. Jenyns and last recorded in 1907 by White (1912). Although still found at Pill, this species appears to have been lost from other coastal sites. It was recorded at Clevedon in 1853 by Leo H. Grindon (White, 1912); in plenty on the hill near the entrenchment, Weston-super-Mare by D.Fry (Murray, 1896) and at Uphill by E.J. Hamlin in 1951 (Sandwith & Sandwith, 1953). The last record from that area of the coast was made by G. Kidder in 1982 at Worlebury Hill, Weston-super-Mare (Roe, 1983). This species was also formerly known from lanes at Shipham and Sidcot; in a field hedge at Stanton Drew, found in 1887 by D. Fry (White, 1912); at Burledge Hill south of Bishop Sutton, found by R.M. Payne and R.G.B. Roe in 1985 (Willis, 1987); in a field by a lane leading from Ursleigh Hill to Publow, found in 1907 by D. Fry (White, 1912); on the border of a wood near Bathampton, found by Miss Peck (White, 1912) and in a lane between Burnt House [Odd Down, Bath] and Priston, found by D. Coombe in 1944. There are no recent records for any of these sites. *Allium oleraceum* has declined throughout its range; sites in Somerset are near the western edge of its distribution in Britain.



*Allium oleraceum* at Lytes Cary (2012). Photo: HJC

## VC5

Ford Farm		SS9140	2010	RJB	
Old Cleeve		ST036415	2002	JW	Along footpath from Old Cleeve to Bilbrook
Cleeve Hill	SSSI	ST052429	1992	IPG	10 plants on edge of scrub by path
Huish		ST28102185	2013	JP	c. 30 plants, W side of hedgerow along footpath
Currymead Lane		ST38542355	2011	JP	9 fruiting spikes, corner of arable headland



Distribution of *Allium oleraceum* in Somerset, mapped using MapMate.  
Black dots are post-2000 records; red squares are 1987-1999 records



*Allium oleraceum* at Pill (2012). Photo: FJR